arXiv:1404.1489v2 [math.CA] 8 Apr 2014

A logarithmic mean and intersections of osculating hyperplanes in R^n

Alan Horwitz
708A Putnam Blvd.
Wallingford, PA 19086-6701
alh4@psu.edu

4/8/14

Abstract

We discuss a special case of the means defined in [1]. Let C be the curve in R^n with vector equation $\hat{\alpha}(t) = \langle t, t\log t, ..., t(\log t)^{n-1}\rangle$. Let $0 < a_1 < \cdots < a_n$ and let O_k be the osculating hyperplane to C at a_k. Then we show that $O_1, ..., O_n$ have a unique point of intersection, $P = (i_1, ..., i_n) \in R^n$, and in particular, i_1 equals the mean $M(a_1, ..., a_n) = (n-1)!\sum_{j=1}^{n} \frac{a_j}{\prod\limits_{\substack{i=1\\ i\neq j}}^{n} (\ln a_j - \ln a_i)}$ = the logarithmic mean of Neuman.

Key Words: logarithmic mean, osculating hyperplane, Wronskian

1 Introduction

For $n \geq 3$, let C be the curve in R^n with vector equation $\hat{\alpha}(t) = \langle x_1(t), ..., x_n(t)\rangle$, let $W_{j,n}(t) = W\left(x_1'(t), ..., x_{j-1}'(t), x_{j+1}'(t), ..., x_n'(t)\right) =$
the Wronskian of $x_1'(t), ..., x_{j-1}'(t), x_{j+1}'(t), ..., x_n'(t), j = 1, ..., n$, let $\hat{x}$ be the vector $\langle x_1, ..., x_n\rangle$, and let $\hat{n}(t)$ be the vector $\langle W_{1,n}(t), -W_{2,n}(t), ..., (-1)^{n+1}W_{n,n}(t)\rangle$. In [1] we defined the **osculating hyperplane**, O_a, to C at $t = a$ to be the hyperplane in R^n with equation

$$\hat{x} \cdot \hat{n}(a) = \hat{\alpha}(a) \cdot \hat{n}(a), \text{ assuming that } \hat{n}(a) \neq \hat{0}.$$

It is not hard to show(see [1]) that O_a has nth order contact with C at $t = a$. That is, if $C_a(t) = (\hat{\alpha}(t) - \hat{\alpha}(a)) \cdot \hat{n}(a)$, then $C_a^{(j)}(a) = 0$ for $j = 0, 1, ..., n-1$. This generalizes the osculating plane in R^3, which has 3rd order contact with C at a. For example, if $\hat{\alpha}(t) = \langle t, t^2, t^3, t^4\rangle$, then $W_{1,4}(t) = W\left(2t, 3t^2, 4t^3\right) = 48t^3, W_{2,4}(t) = W\left(1, 3t^2, 4t^3\right) = 72t^2, W_{3,4}(t) = W\left(1, 2t, 4t^3\right) = 48t, W_{3,4}(t) = W\left(1, 2t, 3t^2\right) = 12$, and $\hat{n}(t) = \langle 48t^3, -72t^2, 48t, -12\rangle$. The equation of the osculating hyperplane at $t = 1$ is $\langle x_1, x_2, x_3, x_4\rangle \cdot \hat{n}(1) = \hat{\alpha}(1) \cdot \hat{n}(1)$ or $4x_1 -$

$6x_2 + 4x_3 - x_4 = 1$. $C_1(t) = \langle t-1, t^2-1, t^3-1, t^4-1\rangle \cdot \langle 48, -72, 48, -12\rangle = -12\left(t^4 - 4t^3 + 6t^2 - 4t + 1\right)$, and it then follows that $C_1(1) = C_1'(1) = C_1''(1) = C_1'''(1) = 0$.

In [1] the author proved the following general result about defining means using intersections of osculating hyperplanes to curves in R^n.

Theorem 1 *Let C be the curve in R^n with vector equation $\hat{\alpha}(t) = \langle x_1(t), ..., x_n(t)\rangle, t \in I = [a, b]$, where each $x_k \in C^{n-1}(I)$ and is strictly monotone on I.*

Let $W_{j,n}(t) = W(x_1'(t), ..., x_{j-1}'(t), x_{j+1}'(t), ..., x_n'(t))$ be the Wronskian of $x_1'(t), ..., x_{j-1}'(t), x_{j+1}'(t), ..., x_n'(t)$. Assume that every subset of $\{W_{1,n}, ..., W_{n,n}\}$ is an extended complete Chebyshev system on I. Let $a = a_1 < \cdots < a_n = b$ be n given points in I, and let O_k be the osculating hyperplane to C at a_k. Then

(1) $O_1, ..., O_n$ have a unique point of intersection, P, in R^n, and

(2) If $P = (i_1, ..., i_n)$, then $a_1 < x_k^{-1}(i_k) < a_n$ for $k = 1, 2, ..., n$

By Theorem 1, one can define n symmetric means in $a_1, ..., a_n$ as follows: $M_k(a_1, ..., a_n) = x_k^{-1}(i_k), k = 1, ..., n$. In particular, we showed in [1] that if $x_k(t) = t^k$, $k = 1, ..., n-2$, $x_{n-1}(t) = \log t$, and $x_n(t) = \dfrac{1}{t}$, then $M_n(a_1, ..., a_n) = P(a_1, ..., a_n)$, where P is the logarithmic mean in n variables defined by Pittenger [5]. At the end of [1] we stated that perhaps another interesting generalization of the logarithmic mean to n variables would be $M_1(a_1, ..., a_n)$, where $x_k(t) = t(\log t)^{k-1}$, $k = 1, ..., n$. We never pursued that, but the point of this paper is to prove that $M_1(a_1, ..., a_n)$ equals the following logarithmic mean in n variables defined by Neuman [4]: $L_N(a_1, ..., a_n) = (n-1)!\displaystyle\sum_{j=1}^{n} \frac{a_j}{\prod\limits_{\substack{i=1\\ i\neq j}}^{n} (\ln a_j - \ln a_i)}$. That is, we show that Neuman's logarithmic mean equals the x coordinate of the intersection of the osculating hyperplanes to the curve $\hat{\alpha}(t) = \langle t, t\log t, ..., t(\log t)^{n-1}\rangle$. L_N was also defined in a different way(and unknowingly) by Xiao and Zhang [7]. Mustonen [3] gives a good summary of these connections and other generalizations. See also the paper by Merikoski [2]. The methods used in this paper are decidedly different than those in the papers just cited. We now state our main result.

Theorem 2 *For $n \geq 3$, let C be the curve in R^n with vector equation $\hat{\alpha}(t) = \langle t, t\log t, ..., t(\log t)^{n-1}\rangle$. Let $0 < a_1 < \cdots < a_n$ and let O_k be the osculating hyperplane to C at a_k. Then $O_1, ..., O_n$ have a unique point of intersection, $P = (i_1, ..., i_n) \in R^n$, and*

$$M_1(a_1, ..., a_n) = (n-1)!\sum_{j=1}^{n} \frac{a_j}{\prod\limits_{\substack{i=1\\ i\neq j}}^{n} (\ln a_j - \ln a_i)}. \tag{1}$$

Note that for $n = 2$, the x coordinate of the point of intersection of the tangent lines to the curve $\hat{\alpha}(t) = \langle t, t\log t\rangle$ is the well known logarithmic mean

$L(a,b) = \frac{b-a}{\ln b - \ln a}$ *in two variables. So in a certain sense the logarithmic mean* $L_N(a_1,...,a_n)$ *above is a natural generalization of the logarithmic mean in two variables since it involves intersections of osculating hyperplanes to a curve in* R^n *whose first two components are* t *and* $t\log t$*, and where the remaining components follow the "natural pattern" of the first two components.*

Remark 1 *Using the curve from Theorem 2 and Theorem 1, one also obtains means* $M_k(a_1,...,a_n) = x_k^{-1}(i_k), k = 2,...,n$*. Of course those means involve the inverse of the function* $y = t(\log t)^k, k \geq 2$*, which is not an elementary function.*

2 Preliminary Material

If $f_1,..,f_n$ are n given functions of t, then we let

$$W(f_1,..,f_n)(t) = \begin{vmatrix} f_1(t) & f_2(t) & \cdots & f_n(t) \\ f'_1(t) & f'_2(t) & \cdots & f'_n(t) \\ \vdots & \vdots & \cdots & \vdots \\ f_1^{(n-1)}(t) & f_2^{(n-1)}(t) & \cdots & f_n^{(n-1)}(t) \end{vmatrix}$$

denote the Wronskian determinant. Throughout the rest of the paper, we define

$$\begin{aligned} x_k(t) &= t(\log t)^{k-1}, k = 1,...,n. \qquad (2) \\ W_{k,n}(t) &= W(x'_1(t),...,x'_{k-1}(t),x'_{k+1}(t),...,x'_n(t)), k = 1,...,n. \end{aligned}$$

Lemma 1 *For* $r \geq 2$

$$x_{k+1}^{(r)}(t) = k\sum_{j=1}^{r-1} \frac{(-1)^{j+1}(j-1)!\binom{r-2}{j-1}}{t^j} x_k^{(r-j)}(t). \qquad (3)$$

Proof. We use induction in r. So suppose that (3) holds for some positive integer $r \geq 2$.

$$x_{k+1}^{(r+1)}(t) = \frac{d}{dt}x_{k+1}^{(r)}(t) = k\frac{d}{dt}\sum_{j=1}^{r-1}\frac{(-1)^{j+1}(j-1)!\binom{r-2}{j-1}}{t^j}x_k^{(r-j)}(t) =$$

$$k\sum_{j=1}^{r-1}(-1)^{j+1}(j-1)!\tbinom{r-2}{j-1}\frac{d}{dt}(t^{-j}x_k^{(r-j)}(t)) =$$

$$k\sum_{j=1}^{r-1}(-1)^{j+1}(j-1)!\tbinom{r-2}{j-1}(t^{-j}x_k^{(r-j+1)}(t) - jt^{-j-1}x_k^{(r-j)}(t)) =$$

$$k\sum_{j=1}^{r-1}\frac{(-1)^{j+1}(j-1)!\binom{r-2}{j-1}}{t^j}x_k^{(r+1-j)}(t) - k\sum_{j=1}^{r-1}\frac{(-1)^{j+1}j!\binom{r-2}{j-1}}{t^{j+1}}x_k^{(r-j)}(t) =$$

$$k\sum_{j=1}^{r-1}\frac{(-1)^{j+1}(j-1)!\binom{r-2}{j-1}}{t^j}x_k^{(r+1-j)}(t) - k\sum_{l=2}^{r}\frac{(-1)^{l}(l-1)!\binom{r-2}{l-2}}{t^l}x_k^{(r+1-l)}(t) =$$

$k\sum_{j=1}^{r-1}\frac{(-1)^{j+1}(j-1)!\binom{r-2}{j-1}}{t^j}x_k^{(r+1-j)}(t)+k\sum_{j=2}^{r}\frac{(-1)^{j+1}(j-1)!\binom{r-2}{j-2}}{t^j}x_k^{(r+1-j)}(t)$.

Using the identity $\binom{r-2}{j-1}+\binom{r-2}{j-2}=\binom{r-1}{j-1}$, we have

$$(j-1)!\binom{r-2}{j-1}+(j-1)!\binom{r-2}{j-2}=(j-1)!\binom{r-1}{j-1},$$

which implies that

$k\sum_{j=1}^{r-1}\frac{(-1)^{j+1}(j-1)!\binom{r-2}{j-1}}{t^j}x_k^{(r+1-j)}(t)+k\sum_{j=2}^{r}\frac{(-1)^{j+1}(j-1)!\binom{r-2}{j-2}}{t^j}x_k^{(r+1-j)}(t)=$

$k\sum_{j=2}^{r-1}\frac{(-1)^{j+1}(j-1)!\binom{r-1}{j-1}}{t^j}x_k^{(r+1-j)}(t)+\frac{k}{t}x_k^{(r)}(t)+\frac{k}{t^r}(-1)^{r+1}(r-1)!x_k^{(1)}(t)=$

$k\sum_{j=1}^{r}\frac{(-1)^{j+1}(j-1)!\binom{r-1}{j-1}}{t^j}x_k^{(r+1-j)}(t)$.

To start the induction, $x'_{k+1}(t)=\frac{k}{t}x_k(t)+\frac{1}{t}x_{k+1}(t)$, which implies that

$x''_{k+1}(t)=\frac{k}{t}x'_k(t)-\frac{k}{t^2}x_k(t)+\frac{1}{t}x'_{k+1}(t)-\frac{1}{t^2}x_{k+1}(t)=\frac{k}{t}x'_k(t)-\frac{k}{t^2}x_k(t)+\frac{1}{t}\left(\frac{k}{t}x_k(t)+\frac{1}{t}x_{k+1}(t)\right)-\frac{1}{t^2}x_{k+1}(t)=$

$\frac{k}{t}x'_k(t)-\frac{k}{t^2}x_k(t)+\frac{k}{t^2}x_k(t)+\frac{1}{t^2}x_{k+1}(t)-\frac{1}{t^2}x_{k+1}(t)=\frac{k}{t}x'_k(t)$, which is (3) with $r=2$. ■

Notation 1 *Let* $a_{r,j}=(-1)^{j+1}(j-1)!\binom{r-2}{j-1}$. *Then Lemma 1 can be written*

$$x_{k+1}^{(r)}(t)=k\sum_{j=1}^{r-1}\frac{a_{r,j}}{t^j}x_k^{(r-j)}(t). \tag{4}$$

Lemma 2 *For* $k\geq 2$, $x_k^{(r)}(1)=0$ *for any* $r\leq k-2$, *and for* $r\geq 2$, $x_r^{(r-1)}(1)=(r-1)!$

Proof. For the first part, we use induction in k. So assume that $x_k^{(l)}(1)=0$ for $l\leq k-2$. By (4), $x_{k+1}^{(r)}(1)=k\sum_{j=1}^{r-1}a_{r,j}x_k^{(r-j)}(1)$. Suppose that $r\leq k-1$. Then $r-j\leq k-j-1\leq k-2$, which implies that $x_k^{(r-j)}(1)=0$; Thus $x_{k+1}^{(r)}(1)=0$ whenever $r\leq(k+1)-2$. To start the induction, consider $x'_k(t)=(\log t)^{k-2}(k-1+\log t)\Rightarrow x'_k(1)=0$ for $k\geq 3$. $x''_{k+1}(t)=\frac{k}{t}x'_k(t)$ for $k\geq 1\Rightarrow x''_k(1)=0$ for $k\geq 4$.

For the second part, we use induction in r. So assume that $x_r^{(r-1)}(1) = (r-1)!$. By (4), $x_{r+1}^{(r)}(1) = r\sum_{j=1}^{r-1} a_{r,j}x_r^{(r-j)}(1) = r\left(a_{r,1}x_r^{(r-1)}(1) + \sum_{j=2}^{r-1} a_{r,j}x_r^{(r-j)}(1)\right) =$ $r(r-1)!$ since $x_r^{(r-j)}(1) = 0$ for $j \geq 2$ by the first part of Lemma 2 just proven. Thus $x_{r+1}^{(r)}(1) = r!$. To start the induction, $x_2'(1) = 1 = 1!$ ■

Lemma 3 $\displaystyle\sum_{k=1}^{n} \frac{(-1)^{k-1}}{(n+1-k)!(k-1)!} = \frac{(-1)^{n+1}}{n!}$

Proof. Follows immediately from the binomial expansion of $1+x$ with $x=-1$ and we omit the details. ■

Lemma 4 $\displaystyle\sum_{k=1}^{n}\left[(-1)^{k-1}\frac{1}{(k-1)!}\left(\sum_{j=0}^{n-k}\frac{x^{n-1-j}}{(n-k-j)!}\right)\right] = 1$ *and*

$\displaystyle\sum_{k=2}^{n}\left[(-1)^{k-1}\frac{1}{(k-2)!}\sum_{j=0}^{n-k}\frac{x^{n-j-2}}{(n-k-j)!}\right] = -1$

Proof. Let $d_n = \displaystyle\sum_{k=1}^{n}\left[(-1)^{k-1}\tfrac{1}{(k-1)!}\sum_{j=0}^{n-k}\tfrac{x^{n-1-j}}{(n-k-j)!}\right]$. Using induction in n, we assume that $d_n = 1$.

$d_{n+1} = \displaystyle\sum_{k=1}^{n+1}\left[(-1)^{k-1}\tfrac{1}{(k-1)!}\left(\sum_{j=0}^{n+1-k}\tfrac{x^{n-j}}{(n+1-k-j)!}\right)\right] =$

$(-1)^n\frac{1}{n!}x^n + \displaystyle\sum_{k=1}^{n}\left[(-1)^{k-1}\tfrac{1}{(k-1)!}\left(\sum_{j=0}^{n+1-k}\tfrac{x^{n-j}}{(n+1-k-j)!}\right)\right] =$

$(-1)^n\frac{1}{n!}x^n + \displaystyle\sum_{k=1}^{n}\left[(-1)^{k-1}\tfrac{1}{(k-1)!}\left(\tfrac{x^n}{(n+1-k)!} + \sum_{j=1}^{n+1-k}\tfrac{x^{n-j}}{(n+1-k-j)!}\right)\right] =$

$(-1)^n\frac{1}{n!}x^n + x^n\displaystyle\sum_{k=1}^{n}\tfrac{(-1)^{k-1}}{(n+1-k)!(k-1)!} + \sum_{k=1}^{n}\left[(-1)^{k-1}\tfrac{1}{(k-1)!}\sum_{j=0}^{n-k}\tfrac{x^{n-1-j}}{(n-k-j)!}\right] =$

$(-1)^n\frac{1}{n!}x^n + x^n\displaystyle\sum_{k=1}^{n}\tfrac{(-1)^{k-1}}{(n+1-k)!(k-1)!} + 1 = (-1)^n\frac{1}{n!}x^n + x^n\frac{(-1)^{n+1}}{n!} + 1$(by Lemma 3) $= 1$.

Since $d_1 = 1$, that completes the proof of the first part of Lemma 4. The proof of the second part of Lemma 4 is similar and we omit it. The second part of Lemma 4 also follows from the first part after some manipulations and Lemma 3. ■

Before proving one of our main results, we introduce the functions $v_{k,n}$ below.

Lemma 5 *For $1 \leq k \leq n$ and $n \geq 3$, let*

$$v_{k,n}(t) = \frac{\prod_{r=0}^{n-1} r!}{(k-1)!} \frac{1}{t^{(n-2)(n-1)/2}} \sum_{j=0}^{n-k} \frac{(\ln t)^{n-k-j}}{(n-k-j)!}, t > 0.$$

Then

$$\begin{aligned} v_{k+1,n+1}(t) &= \frac{n!}{k} \frac{1}{t^{n-1}} v_{k,n}(t) \\ v_{1,n+1}(t) &= \left(\prod_{r=0}^{n-1} r! \right) \frac{(\ln t)^n}{t^{n(n-1)/2}} + \frac{n!}{t^{n-1}} v_{1,n}(t). \end{aligned}$$

Proof. $v_{k+1,n+1}(t) = \frac{\prod_{r=0}^{n-1} r!}{k!} \frac{(\ln t)^{n-k-j}}{t^{n(n-1)/2}} \sum_{j=0}^{n-k} \frac{1}{(n-k-j)!} =$

$\frac{\prod_{r=0}^{n-1} r!}{k!} \frac{(\ln t)^{n-k-j}}{t^{(n-1)(n-2)/2} t^{n-1}} \sum_{j=0}^{n-k} \frac{1}{(n-k-j)!} =$

$\frac{n!}{k} \frac{1}{t^{n-1}} \frac{\prod_{r=0}^{n-1} r!}{(k-1)!} \frac{1}{t^{(n-2)(n-1)/2}} \sum_{j=0}^{n-k} \frac{(\ln t)^{n-k-j}}{(n-k-j)!} = \frac{n!}{k} \frac{1}{t^{n-1}} v_{k,n}(t).$

$v_{1,n}(t) = \left(\prod_{r=0}^{n-1} r! \right) \frac{1}{t^{(n-2)(n-1)/2}} \sum_{j=0}^{n-1} \frac{(\ln t)^{n-1-j}}{(n-1-j)!} \Rightarrow v_{1,n+1}(t) =$

$\left(\prod_{r=0}^{n} r! \right) \frac{1}{t^{n(n-1)/2}} \sum_{j=0}^{n} \frac{(\ln t)^{n-j}}{(n-j)!} =$

$\left(\prod_{r=0}^{n} r! \right) \frac{1}{t^{n(n-1)/2}} \left(\frac{(\ln t)^n}{n!} + \sum_{j=1}^{n} \frac{(\ln t)^{n-j}}{(n-j)!} \right) =$

$\left(\prod_{r=0}^{n} r! \right) \frac{(\ln t)^n}{t^{n(n-1)/2}} \left(\frac{1}{n!} + \sum_{j=0}^{n-1} \frac{(\ln t)^{n-1-j}}{(n-1-j)!} \right) =$

$\left(\prod_{r=0}^{n-1} r! \right) \frac{(\ln t)^n}{t^{n(n-1)/2}} + \left(\prod_{r=0}^{n} r! \right) \frac{1}{t^{n(n-1)/2}} \sum_{j=0}^{n-1} \frac{(\ln t)^{n-1-j}}{(n-1-j)!} =$

$\left(\prod_{r=0}^{n-1} r! \right) \frac{(\ln t)^n}{t^{n(n-1)/2}} + \left(\prod_{r=0}^{n-1} r! \right) \frac{n!}{t^{n-1}} \frac{1}{t^{(n-2)(n-1)/2}} \sum_{j=0}^{n-1} \frac{(\ln t)^{n-1-j}}{(n-1-j)!} =$

$\left(\prod_{r=0}^{n-1} r! \right) \frac{(\ln t)^n}{t^{n(n-1)/2}} + \frac{n!}{t^{n-1}} v_{1,n}(t).$ ■

Proposition 1 *Let $v_{k,n}(t)$ be the functions from Lemma 5, and define the vector functions*

$$\begin{aligned}\hat{\alpha}(t) &= \langle x_1(t), ..., x_n(t)\rangle \\ \hat{v}_n(t) &= \langle v_{1,n}(t), -v_{2,n}(t), ..., (-1)^{n-1}v_{n,n}(t)\rangle.\end{aligned}$$

Then

$$\begin{aligned}\hat{\alpha}(t)\cdot\hat{v}_n(t) &= \frac{\prod_{r=0}^{n-1} r!}{t^{n(n-3)/2}} \\ \hat{\alpha}^{(j)}(t)\cdot\hat{v}_n(t) &= 0 \text{ for } j = 1, ..., n-1.\end{aligned}$$

Remark 2 *$\hat{\alpha}$ depends on n as does $\hat{v}_n$, but we suppress this dependence in our notation for convenience.*

Proof. Case 1: $j = 0$

$\vec{\alpha}(t)\cdot\hat{v}_n(t) = \sum_{k=1}^{n}(-1)^{k-1}x_k(t)v_{k,n}(t) = \sum_{k=1}^{n}(-1)^{k-1}t(\ln t)^{k-1}v_{k,n}(t) =$

$$\left(\prod_{r=0}^{n-1} r!\right)\frac{t}{t^{(n-2)(n-1)/2}}\sum_{k=1}^{n}\left[(-1)^{k-1}\frac{1}{(k-1)!}\left(\sum_{j=0}^{n-k}\frac{(\ln t)^{n-1-j}}{(n-k-j)!}\right)\right] = \frac{\prod_{r=0}^{n-1} r!}{t^{n(n-3)/2}}$$

by Lemma 4 with $x = \ln t$.

Case 2: $j = 1$

$x_k(t) = t(\log t)^{k-1} \Rightarrow x_k'(t) = t(k-1)(\ln t)^{k-2}\frac{1}{t} + (\ln t)^{k-1} \Rightarrow$

$x_k'(t) = (\ln t)^{k-2}(k-1+\ln t)$.

Note that if $k = 1$, $x_k'(t) = 1$ for all t, including $t = 1$.

$\hat{\alpha}'(t)\cdot\hat{v}(t) = \sum_{k=1}^{n}(-1)^{k-1}x_k'(t)v_{k,n}(t) = \sum_{k=1}^{n}(-1)^{k-1}(\ln t)^{k-2}(k-1+\ln t)v_{k,n}(t) =$

$$\frac{\prod_{r=0}^{n-1} r!}{t^{(n-2)(n-1)/2}}\sum_{k=1}^{n}\left[(-1)^{k-1}\frac{1}{(k-1)!}(k-1+\ln t)\left(\sum_{j=0}^{n-k}\frac{(\ln t)^{n-j-2}}{(n-k-j)!}\right)\right] =$$

$$\frac{\prod_{r=0}^{n-1} r!}{t^{(n-2)(n-1)/2}}\times\left(\sum_{k=2}^{n}\left[(-1)^{k-1}\frac{1}{(k-2)!}\sum_{j=0}^{n-k}\frac{(\ln t)^{n-j-2}}{(n-k-j)!}\right] + \sum_{k=1}^{n}\left[(-1)^{k-1}\frac{1}{(k-1)!}\sum_{j=0}^{n-k}\frac{(\ln t)^{n-j-1}}{(n-k-j)!}\right]\right) = 0$$

by Lemma 4 with $x = \ln t$.

Case 3: $2 \le j \le n$(note that j is fixed here)

We use induction in n. So assume that $\hat{\alpha}^{(l)}(t)\cdot\hat{v}_n(t) = \sum_{k=1}^{n}(-1)^{k-1}x_k^{(l)}(t)v_{k,n}(t) = 0$ for all $l = 1, ..., n-1$. We have to show that $\hat{\alpha}^{(j)}(t)\cdot\hat{v}_{n+1}(t) = 0$.

Now $\hat{\alpha}^{(j)}(t)\cdot\hat{v}_{n+1}(t) = \sum_{k=1}^{n+1}(-1)^{k-1}x_k^{(j)}(t)v_{k,n+1}(t) = \sum_{k=0}^{n}(-1)^{k}x_{k+1}^{(j)}(t)v_{k+1,n+1}(t) = x_1^{(j)}(t)v_{1,n+1}(t) + \sum_{k=1}^{n}(-1)^{k}x_{k+1}^{(j)}(t)v_{k+1,n+1}(t)$. Since $x_1^{(j)}(t) = 0$ for $j \geq 2$, we have

$$\hat{\alpha}^{(j)}(t)\cdot\hat{v}_{n+1}(t) = \sum_{k=1}^{n}(-1)^{k}x_{k+1}^{(j)}(t)v_{k+1,n+1}(t). \tag{5}$$

By Lemma 5, for $j \geq 2$, $x_{k+1}^{(j)}(t)v_{k+1,n+1}(t) = k\left(\sum_{i=1}^{j-1}\frac{a_{i,j}}{t^j}x_k^{(j-i)}(t)\right)\frac{n!}{k}\frac{1}{t^{n-1}}v_{k,n}(t) = \frac{n!}{t^{n-1}}\left(\sum_{i=1}^{j-1}\frac{a_{i,j}}{t^j}x_k^{(j-i)}(t)\right)v_{k,n}(t)$.

Then $\sum_{k=1}^{n}(-1)^{k}x_{k+1}^{(j)}(t)v_{k+1,n+1}(t) = \frac{n!}{t^{n-1}}\sum_{k=1}^{n}(-1)^{k}\left(\sum_{i=1}^{j-1}\frac{a_{i,j}}{t^j}x_k^{(j-i)}(t)\right)v_{k,n}(t) =$

$\frac{n!}{t^{n-1}}\sum_{k=1}^{n}(-1)^{k}\left(\sum_{i=1}^{j-1}\frac{a_{i,j}}{t^j}x_k^{(j-i)}(t)v_{k,n}(t)\right) =$

$\frac{n!}{t^{n-1}}\sum_{i=1}^{j-1}\left[\frac{a_{i,j}}{t^j}\left(\sum_{k=1}^{n}(-1)^{k}x_k^{(j-i)}(t)v_{k,n}(t)\right)\right] =$

$\frac{n!}{t^{n-1}}\sum_{i=1}^{j-1}\frac{a_{i,j}}{t^j}\left(\sum_{k=1}^{n}(-1)^{k}x_k^{(j-i)}(t)v_{k,n}(t)\right) =$

$-\frac{n!}{t^{n-1}}\sum_{i=1}^{j-1}\frac{a_{i,j}}{t^j}\left(\sum_{k=1}^{n}(-1)^{k-1}x_k^{(j-i)}(t)v_{k,n}(t)\right) =$

$-\frac{n!}{t^{n-1}}\sum_{i=1}^{j-1}\frac{a_{i,j}}{t^j}\left(\sum_{k=1}^{n}(-1)^{k-1}x_k^{(j-i)}(t)v_{k,n}(t)\right).$

Since $j-i \leq n-1$ for $i \geq 1$, $\sum_{k=1}^{n}(-1)^{k-1}x_k^{(j-i)}(t)v_{k,n}(t) = 0$ by the inductive hypothesis.

Thus $\hat{\alpha}^{(j)}(t)\cdot\hat{v}_{n+1}(t) = 0$. To start the induction, for $n = 1$ we have $\hat{\alpha}(t) = \langle t\rangle \Rightarrow \hat{\alpha}^{(j)}(t) = 0 \Rightarrow \hat{\alpha}^{(j)}(t)\cdot\hat{v}_1(t) = 0$ for $j \geq 2$. ■

Remark 3 *Proposition 1 could perhaps also be proven using properties of hypergeometric functions.*

3 Useful Determinants

Lemma 6 *For $n \geq 3$, $W(x_1, ..., x_n)(t) = \frac{\prod_{r=0}^{n-1} r!}{t^{n(n-3)/2}}, t > 0$, where W denotes the Wronskian.*

Proof. It is easy to show that the n functions $\left\{t(\log t)^{k-1}\right\}_{k=1,...,n}$ satisfy the following nth order Euler DE:

$$t^n\frac{d^ny}{dt^n}+\frac{n^2-3n}{2}t^{n-1}\frac{d^{n-1}y}{dt^{n-1}}+\cdots+a_{n-1}t\frac{dy}{dy}+a_ny=0.$$

By Abel's Identity applied to the interval $(0,\infty)$, $W(x_1,...,x_n)(t)=$

$$C_n\exp\left(-\int\frac{\frac{n^2-3n}{2}t^{n-1}}{t^n}dt\right)=C_n\exp\left(-\frac{n^2-3n}{2}\int\frac{dt}{t}\right)=\frac{C_n}{t^{(n^2-3n)/2}}.$$

We shall let $t=1$ to obtain the precise value $C_n=\prod_{r=0}^{n-1}r!$. $W(x_1,...,x_n)(t)=$

$$\begin{vmatrix} x_1(t) & x_2(t) & \cdots & x_n(t)\\ x_1'(t) & x_2'(t) & \cdots & x_n'(t)\\ x_1''(t) & x_2''(t) & \cdots & x_n''(t)\\ \vdots & \vdots & \cdots & \vdots\\ x_1^{(n-1)}(t) & x_2^{(n-1)}(t) & \cdots & x_n^{(n-1)}(t)\end{vmatrix}=\begin{vmatrix} t & x_2(t) & \cdots & x_n(t)\\ 1 & x_2'(t) & \cdots & x_n'(t)\\ 0 & x_2''(t) & \cdots & x_n''(t)\\ \vdots & \vdots & \cdots & \vdots\\ 0 & x_2^{(n-1)}(t) & \cdots & x_n^{(n-1)}(t)\end{vmatrix}\Rightarrow$$

$$W(x_1,...,x_n)(1)=\begin{vmatrix} 1 & 0 & \cdots & 0\\ 1 & x_2'(1) & \cdots & x_n'(1)\\ 0 & x_2''(1) & \cdots & x_n''(1)\\ \vdots & \vdots & \cdots & \vdots\\ 0 & x_2^{(n-1)}(1) & \cdots & x_n^{(n-1)}(1)\end{vmatrix}=$$

$$\begin{vmatrix} x_2'(1) & \cdots & x_n'(1)\\ x_2''(1) & \cdots & x_n''(1)\\ \vdots & \cdots & \vdots\\ x_2^{(n-1)}(1) & \cdots & x_n^{(n-1)}(1)\end{vmatrix}.$$ The diagonal entries are $x_{r+1}^{(r)}(1), r=1,...,n-1$ and for row i we have $\begin{bmatrix} x_2^{(i)}(1) & \cdots & x_n^{(i)}(1)\end{bmatrix}$.

By Lemma 2, the entries in row i, column $j, j\geq i+2$, are each 0. That shows that the matrix $\begin{bmatrix} x_2'(1) & \cdots & x_n'(1)\\ x_2''(1) & \cdots & x_n''(1)\\ \vdots & \cdots & \vdots\\ x_2^{(n-1)}(1) & \cdots & x_n^{(n-1)}(1)\end{bmatrix}$ is upper triangular, which implies that $\begin{vmatrix} x_2'(1) & \cdots & x_n'(1)\\ x_2''(1) & \cdots & x_n''(1)\\ \vdots & \cdots & \vdots\\ x_2^{(n-1)}(1) & \cdots & x_n^{(n-1)}(1)\end{vmatrix}=\prod_{r=1}^{n-1}x_{r+1}^{(r)}(1)=\prod_{r=0}^{n-1}r!$ by Lemma 2.

■

Remark 4 *One could also prove Lemma 6 by instead finding a formula for the Wronskian of* $\left\{(\log t)^{k-1}\right\}_{k=1}^{n}$ *and using well known properties of Wronskians.*

That would be easier if one did not already have the recursion for $x_{k+1}^{(r)}(t)$. Since we use that recursion elsewhere, it was easier to then prove Lemmas 2 first.

Our next result shows that the Wronskians $W_{k,n}$ are in fact identically equal to the functions $v_{k,n}$ from Lemma 5.

Proposition 2 *For $1 \leq k \leq n$ and $n \geq 3$,*

$$W_{k,n}(t) = \frac{\prod_{r=0}^{n-1} r!}{(k-1)!} \frac{1}{t^{(n-2)(n-1)/2}} \sum_{j=0}^{n-k} \frac{(\ln t)^{n-k-j}}{(n-k-j)!}, t > 0.$$

Proof. Consider the following system of linear equations in the unknown functions $u_1(t), ..., u_n(t)$, where $k_n(t) = \frac{\prod_{r=0}^{n-1} r!}{t^{n(n-3)/2}}$:

$$\begin{aligned} x_1(t)u_1(t) + \cdots x_n(t)u_n(t) &= k_n(t) \\ x_1'(t)u_1(t) + \cdots x_n'(t)u_n(t) &= 0 \\ &\vdots \\ x_1^{(n-1)}(t)u_1(t) + \cdots x_n^{(n-1}(t)u_n(t) &= 0. \end{aligned}$$

The coefficent matrix of this linear system has determinant $W(x_1, ..., x_n)(t)$, which is nonzero by Lemma 6. By Cramer's Rule, the unique solution is given by

$$x_j(t) = \frac{\begin{vmatrix} x_1(t) & \cdots & x_{j-1}(t) & k_n(t) & x_{j+1}(t) & \cdots & x_n(t) \\ x_1'(t) & \cdots & x_{j-1}'(t) & 0 & x_{j+1}'(t) & \cdots & x_n'(t) \\ x_1''(t) & \cdots & x_{j-1}''(t) & 0 & x_{j+1}''(t) & \cdots & x_n''(t) \\ \vdots & \cdots & \vdots & \vdots & \vdots & \cdots & \vdots \\ x_1^{(n-1)}(t) & \cdots & x_{j-1}^{(n-1)}(t) & 0 & x_{j+1}^{(n-1)}(t) & \cdots & x_n^{(n-1)}(t) \end{vmatrix}}{W(x_1,...,x_n)(t)}$$

$$j = 1, ..., n.$$

Expand about column j to obtain $x_j(t) = k_n(t)\frac{(-1)^{j+1}W_{j,n}(t)}{W(x_1,...,x_n)(t)} = (-1)^{j+1}W_{j,n}(t)$. By Proposition 1, $x_j(t) = (-1)^{j+1}v_{j,n}(t)$ also satisfies (7). By uniqueness, $W_{j,n}(t) = v_{j,n}(t), j = 1, ..., n.$ ■

Remark 5 *It follows immediately from Proposition 2, that the $W_{k,n}$ also satisfy the following recursion from Lemma 5 for $n \geq 2$:*

$$W_{k+1,n+1}(t) = \frac{n!}{k}\frac{1}{t^{n-1}}W_{k,n}(t), k \geq 1 \quad (6)$$

$$W_{1,n+1}(t) = \left(\prod_{r=0}^{n-1} r!\right)\frac{(\ln t)^n}{t^{n(n-1)/2}} + \frac{n!}{t^{n-1}}W_{1,n}(t). \quad (7)$$

(6) can also be proven using the determinant definition of the $W_{k,n}$ along with standard properties of determinants. However, we found it difficult to prove (7) this way–hence the introduction of the $v_{k,n}$ functions.

Lemma 7 *For $n \geq 3$,*

$$\sum_{k=1}^{n} \left[(-1)^{k+1} b_k^{n-1} \prod_{1 \leq i < j \leq n;\ i,j \neq k} (b_j - b_i) \right] = (-1)^{n-1} \prod_{1 \leq i < j \leq n} (b_j - b_i).$$

Proof. It is well known that the Vandermonde determinant $\begin{vmatrix} 1 & 1 & \cdots & 1 \\ b_1 & b_2 & \cdots & b_n \\ \vdots & \vdots & \vdots & \vdots \\ b_1^{n-2} & b_2^{n-2} & \cdots & b_n^{n-2} \\ b_1^{n-1} & b_2^{n-1} & \cdots & b_n^{n-1} \end{vmatrix} = \prod_{1 \leq i < j \leq n} (b_j - b_i)$, which implies that $\begin{vmatrix} b_1^{n-1} & b_2^{n-1} & \cdots & b_n^{n-1} \\ b_1^{n-2} & b_2^{n-2} & \cdots & b_n^{n-2} \\ \vdots & \vdots & \vdots & \vdots \\ b_1 & b_2 & \cdots & b_n \\ 1 & 1 & \cdots & 1 \end{vmatrix} = (-1)^{n(n-1)/2} \prod_{1 \leq i < j \leq n} (b_j - b_i)$. By expanding $\begin{vmatrix} b_1^{n-1} & b_2^{n-1} & \cdots & b_n^{n-1} \\ b_1^{n-2} & b_2^{n-2} & \cdots & b_n^{n-2} \\ \vdots & \vdots & \vdots & \vdots \\ b_1 & b_2 & \cdots & b_n \\ 1 & 1 & \cdots & 1 \end{vmatrix}$ along the first row and using induction, one has $(-1)^{(n-1)(n-2)/2} \sum_{k=1}^{n} \left[(-1)^{k+1} b_k^{n-1} \prod_{1 \leq i < j \leq n;\ i,j \neq k} (b_j - b_i) \right] = (-1)^{n(n-1)/2} \prod_{1 \leq i < j \leq n} (b_j - b_i)$ and the lemma follows immediately. ■

Proposition 3 *For $n \geq 3$,* $\begin{vmatrix} W_{1,n}(a_1) & -W_{2,n}(a_1) & \cdots & (-1)^{n+1} W_{n,n}(a_1) \\ W_{1,n}(a_2) & -W_{2,n}(a_2) & \cdots & (-1)^{n+1} W_{n,n}(a_2) \\ \vdots & \vdots & \vdots & \vdots \\ W_{1,n}(a_n) & -W_{2,n}(a_n) & \cdots & (-1)^{n+1} W_{n,n}(a_n) \end{vmatrix} = \left(\prod_{r=0}^{n-1} r! \right)^{n-2} \frac{\prod_{1 \leq i < j \leq n} (\ln a_j - \ln a_i)}{\prod_{j=1}^{n} a_j^{(n-1)(n-2)/2}}.$

Proof. We use induction. So assume that $\begin{vmatrix} W_{1,n-1}(a_1) & -W_{2,n-1}(a_1) & \cdots & (-1)^n W_{n-1,n-1}(a_1) \\ W_{1,n-1}(a_2) & -W_{2,n-1}(a_2) & \cdots & (-1)^n W_{n-1,n-1}(a_2) \\ \vdots & \vdots & \vdots & \vdots \\ W_{1,n-1}(a_{n-1}) & -W_{2,n-1}(a_{n-1}) & \cdots & (-1)^n W_{n-1,n-1}(a_{n-1}) \end{vmatrix} =$

$\left(\prod_{r=0}^{n-2} r!\right)^{n-3} \frac{\prod_{1\le i<j\le n-1} (\ln a_j - \ln a_i)}{\prod_{j=1}^{n-1} a_j^{(n-2)(n-3)/2}}$ for any $0 < a_1 < a_2 < \cdots < a_{n-1}$. Using (6) we have $\begin{vmatrix} W_{1,n}(a_1) & -W_{2,n}(a_1) & \cdots & (-1)^{n+1}W_{n,n}(a_1) \\ W_{1,n}(a_2) & -W_{2,n}(a_2) & \cdots & (-1)^{n+1}W_{n,n}(a_2) \\ \vdots & \vdots & \vdots & \vdots \\ W_{1,n}(a_n) & -W_{2,n}(a_n) & \cdots & (-1)^{n+1}W_{n,n}(a_n) \end{vmatrix} =$

$$((n-1)!)^{n-1} \times \begin{vmatrix} W_{1,n}(a_1) & -\frac{W_{1,n-1}(a_1)}{a_1^{n-2}} & \cdots & \frac{(-1)^k W_{k,n-1}(a_1)}{k a_1^{n-2}} & \cdots & \frac{(-1)^{n+1} W_{n-1,n-1}(a_1)}{(n-1)a_1^{n-2}} \\ W_{1,n}(a_2) & -\frac{W_{1,n-1}(a_2)}{a_2^{n-2}} & \cdots & \frac{(-1)^k W_{k,n-1}(a_2)}{k a_2^{n-2}} & \cdots & \frac{(-1)^{n+1} W_{n-1,n-1}(a_2)}{(n-1)a_2^{n-2}} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ W_{1,n}(a_n) & -\frac{W_{1,n-1}(a_n)}{a_n^{n-2}} & \cdots & \frac{(-1)^k W_{k,n-1}(a_n)}{k a_n^{n-2}} & \cdots & \frac{(-1)^{n+1} W_{n-1,n-1}(a_n)}{(n-1)a_n^{n-2}} \end{vmatrix}.$$

Using (7) yields

$$((n-1)!)^{n-1} \times \begin{vmatrix} \left(\prod_{r=0}^{n-2} r!\right) \frac{(\ln a_1)^{n-1}}{a_1^{(n-1)(n-2)/2}} + \frac{(n-1)! W_{1,n-1}(a_1)}{a_1^{n-2}} & -\frac{W_{1,n-1}(a_1)}{a_1^{n-2}} & \cdots & \frac{(-1)^{n+1} W_{n-1,n-1}(a_1)}{(n-1)a_1^{n-2}} \\ \left(\prod_{r=0}^{n-2} r!\right) \frac{(\ln a_2)^{n-1}}{a_2^{(n-1)(n-2)/2}} + \frac{(n-1)! W_{1,n-1}(a_2)}{a_2^{n-2}} & -\frac{W_{1,n-1}(a_2)}{a_2^{n-2}} & \cdots & \frac{(-1)^{n+1} W_{n-1,n-1}(a_2)}{(n-1)a_2^{n-2}} \\ \vdots & \vdots & \vdots & \vdots \\ \left(\prod_{r=0}^{n-2} r!\right) \frac{(\ln a_n)^{n-1}}{a_n^{(n-1)(n-2)/2}} + \frac{(n-1)! W_{1,n-1}(a_n)}{a_n^{n-2}} & -\frac{W_{1,n-1}(a_n)}{a_n^{n-2}} & \cdots & \frac{(-1)^{n+1} W_{n-1,n-1}(a_n)}{(n-1)a_n^{n-2}} \end{vmatrix}.$$

By adding $(n-1)!$ × Col. 2 to Col.1 we have

$$((n-1)!)^{n-1} \times \begin{vmatrix} \left(\prod_{r=0}^{n-2} r!\right) \frac{(\ln a_1)^{n-1}}{a_1^{(n-1)(n-2)/2}} & -\frac{W_{1,n-1}(a_1)}{a_1^{n-2}} & \cdots & \frac{(-1)^k W_{k,n-1}(a_1)}{k a_1^{n-2}} & \cdots & \frac{(-1)^{n+1} W_{n-1,n-1}(a_1)}{(n-1)a_1^{n-2}} \\ \left(\prod_{r=0}^{n-2} r!\right) \frac{(\ln a_2)^{n-1}}{a_2^{(n-1)(n-2)/2}} & -\frac{W_{1,n-1}(a_2)}{a_2^{n-2}} & \cdots & \frac{(-1)^k W_{k,n-1}(a_2)}{k a_2^{n-2}} & \cdots & \frac{(-1)^{n+1} W_{n-1,n-1}(a_2)}{(n-1)a_2^{n-2}} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \left(\prod_{r=0}^{n-2} r!\right) \frac{(\ln a_n)^{n-1}}{a_n^{(n-1)(n-2)/2}} & -\frac{W_{1,n-1}(a_n)}{a_n^{n-2}} & \cdots & \frac{(-1)^k W_{k,n-1}(a_n)}{k a_n^{n-2}} & \cdots & \frac{(-1)^{n+1} W_{n-1,n-1}(a_n)}{(n-1)a_n^{n-2}} \end{vmatrix}.$$

Factoring out $\prod_{r=0}^{n-2} r!$ from Col. 1, factoring out $\frac{1}{k}$ from Column $k+1, k = 1, .., n-1$, and factoring out $\frac{1}{a_j^{n-2}}$ from row $j, j = 1, ..., n$, yields

$$\frac{((n-1)!)^{n-2}\left(\prod_{r=0}^{n-2} r!\right)}{\prod_{j=1}^{n} a_j^{n-2}} \times$$

$$\begin{vmatrix} \frac{(\ln a_1)^{n-1}}{a_1^{(n-2)(n-3)/2}} & -W_{1,n-1}(a_1) & \cdots & (-1)^k W_{k,n-1}(a_1) & \cdots & (-1)^{n+1} W_{n-1,n-1}(a_1) \\ \frac{(\ln a_2)^{n-1}}{a_2^{(n-2)(n-3)/2}} & -W_{1,n-1}(a_2) & \cdots & (-1)^k W_{k,n-1}(a_2) & \cdots & (-1)^{n+1} W_{n-1,n-1}(a_2) \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \frac{(\ln a_n)^{n-1}}{a_n^{(n-2)(n-3)/2}} & -W_{1,n-1}(a_n) & \cdots & (-1)^k W_{k,n-1}(a_n) & \cdots & (-1)^{n+1} W_{n-1,n-1}(a_n) \end{vmatrix}.$$

By expanding about Col. 1 we obtain

$$\frac{((n-1)!)^{n-2}\left(\prod_{r=0}^{n-2} r!\right)}{\prod_{j=1}^{n} a_j^{n-2}} \times$$

$$\left(\begin{array}{c} \frac{(\ln a_1)^{n-1}}{a_1^{(n-2)(n-3)/2}} \begin{vmatrix} -W_{1,n-1}(a_2) & \cdots & (-1)^{n+1} W_{n-1,n-1}(a_2) \\ \vdots & \vdots & \vdots \\ -W_{1,n-1}(a_n) & \cdots & (-1)^{n+1} W_{n-1,n-1}(a_n) \end{vmatrix} - \\ \frac{(\ln a_2)^{n-1}}{a_2^{(n-2)(n-3)/2}} \begin{vmatrix} -W_{1,n-1}(a_1) & \cdots & (-1)^{n+1} W_{n-1,n-1}(a_1) \\ -W_{1,n-1}(a_3) & \cdots & (-1)^{n+1} W_{n-1,n-1}(a_3) \\ \vdots & \vdots & \vdots \\ -W_{1,n-1}(a_n) & \cdots & (-1)^{n+1} W_{n-1,n-1}(a_n) \end{vmatrix} + \cdots + \\ (-1)^{n+1} \frac{(\ln a_n)^{n-1}}{a_n^{(n-2)(n-3)/2}} \begin{vmatrix} -W_{1,n-1}(a_1) & \cdots & (-1)^{n+1} W_{n-1,n-1}(a_1) \\ \vdots & \vdots & \vdots \\ -W_{1,n-1}(a_{n-1}) & \cdots & (-1)^{n+1} W_{n-1,n-1}(a_{n-1}) \end{vmatrix} \end{array} \right).$$

Factoring out -1 from each column of each determinant and using the induction hypothesis gives

$$(-1)^{n-1} \frac{((n-1)!)^{n-2}\left(\prod_{r=0}^{n-2} r!\right)}{\prod_{j=1}^{n} a_j^{n-2}} \times$$

$$\left(\begin{array}{c} \left(\prod_{r=0}^{n-2} r!\right)^{n-3} \frac{(\ln a_1)^{n-1}}{a_1^{(n-2)(n-3)/2}} \frac{\prod_{2 \le i < j \le n} (\ln a_j - \ln a_i)}{\prod_{j=2}^{n} a_j^{(n-2)(n-3)/2}} + \cdots + \\ (-1)^{n+1} \left(\prod_{r=0}^{n-2} r!\right)^{n-3} \frac{(\ln a_n)^{n-1}}{a_n^{(n-2)(n-3)/2}} \frac{\prod_{1 \le i < j \le n-1} (\ln a_j - \ln a_i)}{\prod_{j=1}^{n-1} a_j^{(n-2)(n-3)/2}} \end{array} \right)$$

$$= (-1)^{n-1}\frac{\left(\prod_{r=0}^{n-1} r!\right)^{n-2}}{\prod_{j=1}^{n} a_j^{(n-1)(n-2)/2}} \times$$
$$((\ln a_1)^{n-1} \prod_{2\le i<j\le n} (\ln a_j - \ln a_i) + \cdots$$
$$+(-1)^{n+1} (\ln a_n)^{n-1} \prod_{1\le i<j\le n-1} (\ln a_j - \ln a_i)).$$

Applying Lemma 7 to each term of the sum in parentheses yields $\frac{\left(\prod_{r=0}^{n-1} r!\right)^{n-2}}{\prod_{j=1}^{n} a_j^{(n-1)(n-2)/2}} \prod_{1\le i<j\le n} (\ln a_j - \ln a_i)$. For $n = 3$ we have $W_{1,3}(t) = \frac{\ln^2 t + 2\ln t + 2}{t}$, $W_{2,3}(t) = 2\frac{\ln t + 1}{t}$, and $W_{3,3}(t) = \frac{1}{t}$. Thus $\begin{vmatrix} W_{1,3}(a_1) & -W_{2,3}(a_1) & W_{3,3}(a_1) \\ W_{1,3}(a_2) & -W_{2,3}(a_2) & W_{3,3}(a_2) \\ W_{1,3}(a_3) & W_{2,3}(a_3) & W_{3,3}(a_3) \end{vmatrix} =$

$\begin{vmatrix} \frac{\ln^2 a_1 + 2\ln a_1 + 2}{a_1} & -2\frac{\ln a_1 + 1}{a_1} & \frac{1}{a_1} \\ \frac{\ln^2 a_2 + 2\ln a_2 + 2}{a_2} & -2\frac{\ln a_2 + 1}{a_2} & \frac{1}{a_2} \\ \frac{\ln^2 a_3 + 2\ln a_3 + 2}{a_3} & -2\frac{\ln a_3 + 1}{a_3} & \frac{1}{a_3} \end{vmatrix} = \frac{2(\ln a_3 - \ln a_1)(\ln a_3 - \ln a_2)(\ln a_2 - \ln a_1)}{a_1 a_2 a_3}$ (after some simplification), which equals $\left(\prod_{r=0}^{n-1} r!\right)^{n-2} \frac{\prod_{1\le i<j\le n} (\ln a_j - \ln a_i)}{\prod_{j=1}^{n} a_j^{(n-1)(n-2)/2}}$ for $n = 3$. ■

Proposition 4 *For $n \ge 3$, let $k_n(t) = \dfrac{\prod_{r=0}^{n-1} r!}{t^{n(n-3)/2}}$. Then*

$$\begin{vmatrix} k_n(a_1) & -W_{2,n}(a_1) & \cdots & (-1)^{n+1} W_{n,n}(a_1) \\ k_n(a_2) & -W_{2,n}(a_2) & \cdots & (-1)^{n+1} W_{n,n}(a_2) \\ \vdots & \vdots & \vdots & \vdots \\ k_n(a_n) & -W_{2,n}(a_n) & \cdots & (-1)^{n+1} W_{n,n}(a_n) \end{vmatrix} =$$
$$(-1)^{n-1}(n-1)! \left(\prod_{r=0}^{n-1} r!\right)^{n-2} \frac{\sum_{i=1}^{n} \left(\prod_{\substack{1\le j<k\le n \\ j\ne i, k\ne i}} (-1)^{i+1} a_i (\ln a_k - \ln a_j) \right)}{\prod_{j=1}^{n} a_j^{(n-1)(n-2)/2}}.$$

Proof. We again use induction. So assume that

$$\begin{vmatrix} k_{n-1}(a_1) & -W_{2,n-1}(a_1) & \cdots & (-1)^{n} W_{n-1,n-1}(a_1) \\ k_{n-1}(a_2) & -W_{2,n-1}(a_2) & \cdots & (-1)^{n} W_{n-1,n-1}(a_2) \\ \vdots & \vdots & \vdots & \vdots \\ k_{n-1}(a_{n-1}) & -W_{2,n-1}(a_{n-1}) & \cdots & (-1)^{n} W_{n-1,n-1}(a_{n-1}) \end{vmatrix} =$$

$(-1)^n(n-2)!\left(\prod_{r=0}^{n-2} r!\right)^{n-3} \dfrac{\sum_{i=1}^{n-1}\left(\prod_{\substack{1\le j<k\le n-1\\ j\neq i,k\neq i}} (-1)^{i+1}a_i(\ln a_k-\ln a_j)\right)}{\prod_{m=1}^{n-1} a_j^{(n-2)(n-3)/2}}$ for any $0<a_1<a_2<\cdots<a_{n-1}$. Using (6) we have

$$\begin{vmatrix} k_n(a_1) & -W_{2,n}(a_1) & \cdots & (-1)^{n+1}W_{n,n}(a_1)\\ k_n(a_2) & -W_{2,n}(a_2) & \cdots & (-1)^{n+1}W_{n,n}(a_2)\\ \vdots & \vdots & \vdots & \vdots\\ k_n(a_n) & -W_{2,n}(a_n) & \cdots & (-1)^{n+1}W_{n,n}(a_n)\end{vmatrix} =$$

$$\begin{vmatrix} \frac{\prod_{r=0}^{n-1} r!}{a_1^{n(n-3)/2}} & -\frac{(n-1)!W_{1,n-1}(a_1}{a_1^{n-2}}) & \cdots & \frac{(-1)^{n+1}(n-1)!W_{n-1,n-1}(a_1)}{a_1^{n-2}(n-1)}\\ \frac{\prod_{r=0}^{n-1} r!}{a_2^{n(n-3)/2}} & -\frac{(n-1)!W_{1,n-1}(a_2)}{a_2^{n-2}} & \cdots & \frac{(-1)^{n+1}(n-1)!W_{n-1,n-1}(a_2)}{a_2^{n-2}(n-1)}\\ \vdots & \vdots & \vdots & \vdots\\ \frac{\prod_{r=0}^{n-1} r!}{a_n^{n(n-3)/2}} & -\frac{(n-1)!W_{1,n-1}(a_n)}{a_n^{n-2}} & \cdots & \frac{(-1)^{n+1}(n-1)!W_{n-1,n-1}(a_n)}{a_n^{n-2}(n-1)}\end{vmatrix}.$$

Factoring out $\prod_{r=0}^{n-1} r!$ from Col. 1, factoring out $\frac{(n-1)!}{k}$ from Column $k+1, k=1,..,n-1$, and factoring out $\frac{1}{a_j^{n-2}}$ from row $j, j=1,...,n$, we obtain

$$\frac{((n-1)!)^{n-2}\prod_{r=0}^{n-1} r!}{\prod_{j=1}^{n} a_j^{n-2}}\begin{vmatrix} \frac{1}{a_1^{(n-1)(n-4)/2}} & -W_{1,n-1}(a_1) & \cdots & (-1)^{n+1}W_{n-1,n-1}(a_1)\\ \frac{1}{a_2^{(n-1)(n-4)/2}} & -W_{1,n-1}(a_2) & \cdots & (-1)^{n+1}W_{n-1,n-1}(a_2)\\ \vdots & \vdots & \vdots & \vdots\\ \frac{1}{a_n^{(n-1)(n-4)/2}} & -W_{1,n-1}(a_n) & \cdots & (-1)^{n+1}W_{n-1,n-1}(a_n)\end{vmatrix}.$$

Expanding about Col. 1 gives

$$\frac{((n-1)!)^{n-2}\prod_{r=0}^{n-1} r!}{\prod_{j=1}^{n} a_j^{n-2}} \times$$

$$\left(\begin{array}{c} \frac{1}{a_1^{(n-1)(n-4)/2}} \begin{vmatrix} -W_{1,n-1}(a_2) & \cdots & (-1)^{n+1}W_{n-1,n-1}(a_2) \\ \vdots & \vdots & \vdots \\ -W_{1,n-1}(a_n) & \cdots & (-1)^{n+1}W_{n-1,n-1}(a_n) \end{vmatrix} - \\ \frac{1}{a_2^{(n-1)(n-4)/2}} \begin{vmatrix} -W_{1,n-1}(a_1) & \cdots & (-1)^{n+1}W_{n-1,n-1}(a_1) \\ -W_{1,n-1}(a_3) & \cdots & (-1)^{n+1}W_{n-1,n-1}(a_3) \\ \vdots & \vdots & \vdots \\ -W_{1,n-1}(a_n) & \cdots & (-1)^{n+1}W_{n-1,n-1}(a_n) \end{vmatrix} + \cdots \\ +(-1)^{n+1}\frac{1}{a_n^{(n-1)(n-4)/2}} \begin{vmatrix} -W_{1,n-1}(a_1) & \cdots & (-1)^{n+1}W_{n-1,n-1}(a_1) \\ \vdots & \vdots & \vdots \\ -W_{1,n-1}(a_{n-1}) & \cdots & (-1)^{n+1}W_{n-1,n-1}(a_{n-1}) \end{vmatrix}. \end{array} \right)$$

.

Factoring out -1 from each column of each determinant and using Proposition 3 yields

$$(-1)^{n-1}\frac{((n-1)!)^{n-2}\prod_{r=0}^{n-1} r!}{\prod_{j=1}^{n} a_j^{n-2}} \times$$

$$\left(\begin{array}{c} \frac{1}{a_1^{(n-1)(n-4)/2}} \left(\prod_{r=0}^{n-2} r!\right)^{n-3} \frac{\prod_{2\le i<j\le n}(\ln a_j - \ln a_i)}{\prod_{j=2}^{n} a_j^{(n-2)(n-3)/2}} + \cdots + \\ (-1)^{n+1}\frac{1}{a_n^{(n-1)(n-4)/2}} \left(\prod_{r=0}^{n-2} r!\right)^{n-3} \frac{\prod_{1\le i<j\le n-1}(\ln a_j - \ln a_i)}{\prod_{j=1}^{n-1} a_j^{(n-2)(n-3)/2}} \end{array} \right)$$

=

$$= (-1)^{n-1}\frac{((n-1)!)^{n-2}\left(\prod_{r=0}^{n-1} r!\right)\left(\prod_{r=0}^{n-2} r!\right)^{n-3}}{\prod_{j=1}^{n} a_j^{n-2}} \times$$

$$\left(\frac{a_1 \prod_{2\le i<j\le n}(\ln a_j - \ln a_i)}{\prod_{j=1}^{n} a_j^{(n-2)(n-3)/2}} + \cdots + (-1)^{n+1}\frac{a_n \prod_{1\le i<j\le n-1}(\ln a_j - \ln a_i)}{\prod_{j=1}^{n} a_j^{(n-2)(n-3)/2}} \right)$$

$= (-1)^{n-1}(n-1)!\left(\prod_{r=0}^{n-1} r!\right)^{n-2} \frac{\sum_{i=1}^{n}\left(\prod_{\substack{1\le j<k\le n \\ j\neq i,k\neq i}} (-1)^{i+1}a_i(\ln a_k - \ln a_j)\right)}{\prod_{j=1}^{n} a_j^{(n-1)(n-2)/2}}$. For $n=3$ we have $\begin{vmatrix} k_3(a_1) & -W_{2,3}(a_1) & W_{3,3}(a_1) \\ k_3(a_2) & -W_{2,3}(a_2) & W_{3,3}(a_2) \\ k_3(a_3) & W_{2,3}(a_3) & W_{3,3}(a_3) \end{vmatrix} =$

$\begin{vmatrix} 2 & -2\frac{\ln a_1+1}{a_1} & \frac{1}{a_1} \\ 2 & -2\frac{\ln a_2+1}{a_2} & \frac{1}{a_2} \\ 2 & -2\frac{\ln a_3+1}{a_3} & \frac{1}{a_3} \end{vmatrix} = 4\frac{a_1(\ln a_3-\ln a_2)-a_2(\ln a_3-\ln a_1)+a_3(\ln a_2-\ln a_1)}{a_1a_2a_3}$ (after some simplification), which equals

$(-1)^{n-1}(n-1)!\left(\prod_{r=0}^{n-1} r!\right)^{n-2} \frac{\sum_{i=1}^{n}\left(\prod_{\substack{1\le j<k\le n \\ j\neq i,k\neq i}} (-1)^{i+1}a_i(\ln a_k - \ln a_j)\right)}{\prod_{j=1}^{n} a_j^{(n-1)(n-2)/2}}$ for $n=3$. ■

4 Proof of Theorem 2

Proof. The equation of the osculating hyperplane, O_a, to C at $t=a$ is $\langle x_1,...,x_n\rangle \cdot \hat{n}(a) = \hat{\alpha}(a)\cdot\hat{n}(a)$, where $\hat{\alpha}(t) = \langle t, t\log t, ..., t(\log t)^{n-1}\rangle$,

$\hat{n}(t) = \langle W_{1,n}(t), -W_{2,n}(t), ..., (-1)^{n+1}W_{n,n}(t)\rangle$, $W_{j,n}(t)$ = the Wronskian of $x_1'(t),...,x_{j-1}'(t),x_{j+1}'(t),...,x_n'(t), j=1,...,n$. Thus any intersection point of $O_1,...,O_n$ must be a solution of the linear system $\langle x_1,...,x_n\rangle\cdot\hat{n}(a_j) = \hat{\alpha}(a_j)\cdot\hat{n}(a_j) = 0, j=1,...,n$, which can be written in the form

$$\begin{bmatrix} W_{1,n}(a_1) & -W_{2,n}(a_1) & \cdots & (-1)^{n+1}W_{n,n}(a_1) \\ W_{1,n}(a_2) & -W_{2,n}(a_2) & \cdots & (-1)^{n+1}W_{n,n}(a_2) \\ \vdots & \vdots & \vdots & \vdots \\ W_{1,n}(a_n) & -W_{2,n}(a_n) & \cdots & (-1)^{n+1}W_{n,n}(a_n) \end{bmatrix} \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} = \tag{8}$$
$$\begin{bmatrix} k(a_1) \\ \vdots \\ k(a_n) \end{bmatrix},$$

where $k(t) = \hat{\alpha}(t)\cdot\hat{n}(t) = W(x_1,...,x_n)(t) = \frac{\prod_{r=0}^{n-1} r!}{t^{n(n-3)/2}}$ by Lemma 6. (8) has a

unique solution, $x_1, ..., x_n$, by Proposition 3. By Cramer's Rule,

$$x_1 = \begin{vmatrix} W_{1,n}(a_1) & -W_{2,n}(a_1) & \cdots & (-1)^{n+1}W_{n,n}(a_1) \\ W_{1,n}(a_2) & -W_{2,n}(a_2) & \cdots & (-1)^{n+1}W_{n,n}(a_2) \\ \vdots & \vdots & \vdots & \vdots \\ W_{1,n}(a_n) & -W_{2,n}(a_n) & \cdots & (-1)^{n+1}W_{n,n}(a_n) \end{vmatrix} / \begin{vmatrix} k_n(a_1) & -W_2(a_1) & \cdots & (-1)^{n+1}W_n(a_1) \\ k_n(a_2) & -W_2(a_2) & \cdots & (-1)^{n+1}W_n(a_2) \\ \vdots & \vdots & \vdots & \vdots \\ k_n(a_n) & -W_2(a_n) & \cdots & (-1)^{n+1}W_n(a_n) \end{vmatrix}$$

$$= \frac{(-1)^{n-1}(n-1)!\left(\prod_{r=0}^{n-1} r!\right)^{n-2} \frac{\sum_{i=1}^{n}\left(\prod_{\substack{1\le j<k\le n \\ j\neq i, k\neq i}} (-1)^{i+1}a_i(\ln a_k - \ln a_j)\right)}{\prod_{j=1}^{n} a_j^{(n-1)(n-2)/2}}}{\left(\prod_{r=0}^{n-1} r!\right)^{n-2} \frac{\prod_{1\le i<j\le n}(\ln a_j - \ln a_i)}{\prod_{j=1}^{n} a_j^{(n-1)(n-2)/2}}} \text{ by Propositions 3}$$

and 4. Simplifying gives $\frac{(n-1)!\sum_{i=1}^{n}\left(\prod_{\substack{1\le j<k\le n \\ j\neq i, k\neq i}} (-1)^{n+i}a_i(\ln a_k - \ln a_j)\right)}{\prod_{1\le i<j\le n}(\ln a_j - \ln a_i)}$. By getting a common denominator in the right hand side of (1), it then follows easily that the latter expression equals $(n-1)!\sum_{j=1}^{n} \frac{a_j}{\prod_{\substack{i=1 \\ i\neq j}}^{n}(\ln a_j - \ln a_i)}$. ■

Remark 6 *In [7] the following extension of the Identric mean* $I(a,b) = \left(\frac{a^a}{b^b}\right)^{1/(a-b)}/e$ *to n variables was given:*

$$I_Z(a_1, ..., a_n) = \exp\left[\frac{1}{V(a)}\sum_{i=1}^{n}(-1)^{n+i}a_i^{n-1}V_i(a)\ln a_i - m\right], \text{ where } V(a_1, ..., a_n) = \prod_{1\le j<i\le n}(a_i - a_j), V_i(a_1, ..., a_n) = \begin{vmatrix} 1 & 1 & ... & 1 & 1 & ... & 1 \\ a_1 & a_2 & ... & a_{i-1} & a_{i+1} & ... & a_n \\ a_1^2 & a_2^2 & ... & a_{i-1}^2 & a_{i+1}^2 & ... & a_n^2 \\ ... & ... & ... & ... & ... & ... & ... \\ a_1^{n-2} & a_2^{n-2} & ... & a_{i-1}^{n-2} & a_{i+1}^{n-2} & ... & a_n^{n-2} \end{vmatrix},$$

and $m = \sum_{k=1}^{n-1}\frac{1}{k}$. *For* $n = 3$, *if one lets* $x_1(t) = t$, $x_2(t) = t^2$, $x_3(t) = \log t$, *then*

$M_z(a,b,c) = I_Z(a,b,c) = U_2(a,b,c)$, where U_2 is given in [6]. This probably holds for all n.

Conjecture 1 *If $x_1(t) = t$, $x_2(t) = t^2$, ..., $x_{n-1}(t) = t^{n-1}, x_n(t) = \log t$, then $M_n(a_1, ..., a_n) = I_Z(a_1, ..., a_n)$.*

This conjecture is probably somewhat easier to prove than Theorem 2.

References

[1] A. Horwitz, "Means, Generalized Divided Differences, and Intersections of Osculating Hyperplanes", JMAA 200(1996), 126–148.

[2] Jorma K. Merikoski, "Extending Means Of Two Variables To Several Variables, JIPAM, Volume 5, Issue 3, Article 65, 2004.

[3] S. Mustonen, "Logarithmic mean for several arguments", (2002). ONLINE [http://www.survo.fi/papers/logmean.pdf].

[4] Edward Neuman, "The Weighted Logarithmic Mean", JMAA 188, 885–900(1994).

[5] A. O. Pittenger, The logarithmic mean in n variables, Amer. Math. Monthly 92(1985), 99–104.

[6] K. Stolarsky, "Generalizations of the logarithmic mean", Math. Mag. 48 (1975), 87–92.

[7] Zhen-Gang Xiao and Zhi-Hua Zhang, "The Inequalities $G \leq L \leq I \leq A$ in n Variables", JIPAM, Volume 4, Issue 2, Article 39, 2003.